

Vitamin B Complex

COMPOSITION:

Each 5 ml Adcovit ® syrup contains:

- Thiamine Hydrochloride BP 5 mg (Vit.B1)
- Riboflavin (as Riboflavin-5 Phosphate Sodium BP) 2 mg (Vit.B2)
- Pyridoxine Hydrochloride BP 2 mg (Vit.B_s)
- · Nicotinamide BP 20 mg

PHARMACOLOGY:

Adcovit ® syrup contains the most important members of the vitamin B group in therapeutically balanced proportions. The members of the vitamin B group contained in Adcovit syrup are components of enzyme systems that regulate various stages of carbohydrate, fat and protein metabolism. Each of the components playing a specific biological role.

INDICATION:

Adcovit [®] syrup is indicated for nutritional supplementation as a prophylactic and therapeutic agent against specific vitamin B deficiencies, either single or multiple. Adcovit [®] syrup is also taken along with antibiotic or antitubercular drugs.

DOSE & ADMINISTRATION:

Adcovit [®] Syrup 2-3 teaspoonful daily or as directed by the physician.

CONTRA-INDICATION:

Adcovit ® is contraindicated in patients hypersensitive to any of its components.

WARNING AND PRECAUTION:

Adcovit ® should be given coutiously to patients taking levodopa as phyridoxine reduces the effect of Levodopa.

SIDE EFFECTS:

Multivitamin preparation with ordinary doses of each component is usually nontoxic.

USE IN PREGNANCY & LACTATION:

It is safe to use Adcovit ® Syrup in pregnancy and lactation.

USE IN CHILDREN & ADOLESCENTS:

It is safe to use Adcovit ® Syrup in children & adolescents.

DRUG INTERACTION:

There are no known drug interactions and none well documented.

OVERDOSE:

Taking more medication will not improve the symptoms, rather it may cause poisoining or serious side effects.

STORAGE:

Store below 30° C. Protected from light & moisture. Keep out of the reach of children.

PACKING:

Adcovit [®] Syrup: Each bottle containing 200 ml syrup with a measuring cup.

